

HOMOGENEOUS FOURIER MULTIPLIERS OF MARCINKIEWICZ TYPE

ANTHONY CARBERY AND ANDREAS SEEGER

1. Introduction

Let $m \in L^\infty(\mathbb{R}^2)$ be homogeneous of degree zero. Then m is almost everywhere determined by $h_\pm(\xi_1) = m(\xi_1, \pm 1)$. For $k \in \mathbb{Z}$ let $I_k = [2^{-k-1}, 2^{-k}] \cup [-2^{-k}, -2^{-k-1}]$ and let h_+ and h_- satisfy the condition

$$(1.1) \quad \sup_{k \in \mathbb{Z}} \left(\int_{I_k} |sh'_\pm(s)|^r \frac{ds}{s} \right)^{1/r} < \infty.$$

Rubio de Francia posed the question whether a condition like (1.1) is sufficient to prove that m is a Fourier multiplier of $L^p(\mathbb{R}^2)$, $1 < p < \infty$. An application of the Marcinkiewicz multiplier theorem with L^2 -Sobolev hypotheses (cf. (1.3) and (1.5) below) and interpolation arguments already show that the answer is yes, provided $r > 2$. Recently, Duoandikoetxea and Moyua [15] have shown that the same conclusion can be reached if $r = 2$. On the other hand, since characteristic functions of halfspaces are Fourier multipliers of L^p , $1 < p < \infty$, a simple averaging argument shows that the condition $h' \in L^1$ implies L^p -boundedness for $1 < p < \infty$. Our first theorem shows that the weaker assumption (1.1) with $r = 1$ implies boundedness in $L^p(\mathbb{R}^2)$, for $1 < p < \infty$.

Theorem 1.1. *Suppose that h_+ and h_- satisfy the hypotheses of the Marcinkiewicz multiplier theorem on the real line, that is*

$$(1.2) \quad \sup_{k \in \mathbb{Z}} \int_{I_k} |dh_\pm(s)| \leq A$$

for $I_k = [2^{-k-1}, 2^{-k}] \cup [-2^{-k}, -2^{-k-1}]$. Let $m \in L^\infty(\mathbb{R}^2)$ be homogeneous of degree zero, such that for $\xi_1 \in \mathbb{R}$, $m(\xi_1, 1) = h_+(\xi_1)$ and $m(\xi_1, -1) = h_-(\xi_1)$. Then m is a Fourier multiplier of $L^p(\mathbb{R}^2)$, $1 < p < \infty$, with norm $\leq C A$.

One can obtain a stronger result for fixed $p > 1$ using the space V^q of functions of bounded q -variation. Given an interval I on the real line a function h belongs to $V^q(I)$ if for each partition $\{x_0 < x_1 < \dots < x_N\}$ of I the sum $\sum_{\nu=1}^N |h(x_\nu) - h(x_{\nu-1})|^q$ is bounded and the upper bound of such sums is finite. We denote by $\|h\|_{V^q}^q$ the least upper bound. Then the following result is an immediate consequence of Theorem 1.1 and the interpolation argument in [8].

1991 *Mathematics Subject Classification.* 42B15.

Key words and phrases. Homogeneous multipliers, Differentiation in lacunary directions, Kakeya maximal functions, Multiparameter Calderón-Zygmund theory.

The second author was supported in part by a grant from the National Science Foundation (USA)

Corollary 1.2. *Let m , $h_{\pm}$ and I_k be as above and suppose that*

$$\|h_{\pm}\|_{\infty} + \sup_k \|h_{\pm}\|_{V^q(I_k)} < \infty.$$

Then m is a Fourier multiplier of $L^p(\mathbb{R}^2)$, if $|1/p - 1/2| < 1/2q$.

A slightly weaker result can be formulated in terms of Sobolev spaces. Let β be an even C^∞ function on the real line, supported in $(5/8, 8/5) \cup (-8/5, -5/8)$ and positive in $(1/\sqrt{2}, \sqrt{2}) \cup (-\sqrt{2}, -1/\sqrt{2})$; we shall usually assume that $\sum_{k \in \mathbb{Z}} \beta^2(2^k s) = 1$ for $s \neq 0$. Let $L_\alpha^q(\mathbb{R}^d)$ denote the standard Sobolev space with norm $\|h\|_{L_\alpha^q} = \|\mathcal{F}^{-1}[(1 + |\xi|^2)^{\alpha/2} \hat{h}]\|_q$. Then $L_\alpha^q(\mathbb{R}) \subset V_q$ if $\alpha > 1/q$ and therefore we obtain

Corollary 1.3. *Let $m \in L^\infty(\mathbb{R}^2)$ be homogeneous of degree zero and $h_{\pm}(\xi_1) = m(\xi_1, \pm 1)$. Suppose that $q > 1$ and that*

$$(1.3) \quad \sup_{t \in \mathbb{R}_+} \|\beta h_{\pm}(t \cdot)\|_{L_\alpha^q(\mathbb{R})} < \infty, \quad \alpha > \frac{1}{q}.$$

Then m is a Fourier multiplier of $L^p(\mathbb{R}^2)$ if $|1/p - 1/2| < 1/2q$.

We now compare these results with more standard multiparameter versions of the Hörmander-Marcinkiewicz multiplier theorem. In order to formulate them let

$$\mathcal{D}_j^\alpha g = \mathcal{F}^{-1}[(1 + |\xi_j|^2)^{\alpha/2} \mathcal{F}g]$$

and, for $1 < q < \infty$, let $\mathcal{H}_\alpha^q(\mathbb{R}^n)$ be the *multiparameter* Sobolev space of all functions g , such that

$$\|g\|_{\mathcal{H}_\alpha^q(\mathbb{R}^n)} := \|\mathcal{D}_1^\alpha \dots \mathcal{D}_n^\alpha g\|_{L^q(\mathbb{R}^n)} < \infty.$$

Let β be as above and denote by $\beta_{(i)}$ a copy of β as a function of the ξ_i -variable. Then if $q \geq 2$ the condition

$$(1.4) \quad \sup_{t \in (\mathbb{R}_+)^d} \|\beta_{(1)} \otimes \dots \otimes \beta_{(d)} m(t_1 \cdot, \dots, t_d \cdot)\|_{\mathcal{H}_\alpha^q(\mathbb{R}^d)} < \infty, \quad \alpha > \frac{1}{q}$$

implies that m is a Fourier multiplier of L^p for $|1/p - 1/2| < 1/q$. For $q = 2$ the proof of this result is a variant of Stein's proof of the Hörmander multiplier theorem (see [25, ch.IV]) and the general case follows by an interpolation argument as in [9]. If we apply this result to homogeneous multipliers and set

$$(1.5) \quad m(\xi', \pm 1) = g_{\pm}(\xi'), \quad \xi' \in \mathbb{R}^{d-1}$$

we obtain by a straightforward computation

Corollary 1.4. *Suppose that $r \geq 2$,*

$$(1.6) \quad \sup_{t \in (\mathbb{R}_+)^{d-1}} \|\mathcal{D}_1^{2\gamma} \mathcal{D}_2^\gamma \dots \mathcal{D}_{d-1}^\gamma [\beta_{(1)} \otimes \dots \otimes \beta_{(d-1)} g_{\pm}(t_1 \cdot, \dots, t_{d-1} \cdot)]\|_{L^r(\mathbb{R}^{d-1})} < \infty, \quad \gamma > \frac{1}{r},$$

and that the condition analogous to (1.6) holds for all permutations of the $(s_1, \dots, s_{d-1})$ -variables. Let m be homogeneous of degree zero and related to $g_{\pm}$ by (1.5). Then m is a Fourier multiplier of $L^p(\mathbb{R}^d)$ if $|1/p - 1/2| < 1/r$.

In two dimensions Corollary 1.4 says that if $\alpha > 1/q$, $q \geq 1$, and $\beta g_{\pm}(t \cdot) \in \mathcal{H}_\alpha^{2q}(\mathbb{R})$, uniformly in $t > 0$, then m is a Fourier multiplier of L^p if $|1/p - 1/2| < 1/2q$. Corollary 1.3 is stronger since a compactly supported function in $H_\alpha^{2q}(\mathbb{R})$ belongs to $H_\alpha^q(\mathbb{R})$.

We are now going to discuss variants of Theorem 1.1 in higher dimensions. First if $g_{\pm} \in \mathcal{H}_{\alpha}^q(\mathbb{R}^{d-1})$, $\alpha > 1/q$ and if $g_{\pm}$ are compactly supported in $[1/2, 2]^{d-1}$ then the homogeneous extension m is a Fourier multiplier of $L^p(\mathbb{R}^d)$ if $|1/p - 1/2| < 1/2q$. In fact by a simple averaging argument one sees that the condition $g_{\pm} \in \mathcal{H}_{1+\epsilon}^1$ implies that m is an L^1 multiplier and the general case follows by interpolation. We remark that if $\alpha < |2/p - 1|$ the condition $g_{\pm} \in \mathcal{H}_{\alpha}^q$ (any q) does not imply that m is a Fourier multiplier of L^p . Relevant counterexamples have been pointed out by López-Melero [22] and Christ [7].

Perhaps surprisingly, the situation in higher dimensions changes if one imposes dilation invariant conditions as in Theorem 1.1. One might want to just replace hypothesis (1.2) by the hypotheses of the Marcinkiewicz multiplier theorem in $\mathbb{R}^{d-1}$ ([25, p.108]). However this assumption is not sufficient to deduce that m is a Fourier multiplier of L^p for any $p \neq 2$ (see §3 for the counterexample involving the Kakeya set). However we do have

Theorem 1.5. *Let $m \in L^{\infty}(\mathbb{R}^d)$, $d \geq 2$, be homogeneous of degree zero and let $g_{\pm}$ be as in (1.5). Suppose that $q \geq 2$, and*

$$(1.7) \quad \sup_{t \in (\mathbb{R}_+)^{d-1}} \|\beta_{(1)} \otimes \cdots \otimes \beta_{(d-1)} g_{\pm}(t_1, \dots, t_{d-1})\|_{\mathcal{H}_{\alpha}^q(\mathbb{R}^{d-1})} < \infty, \quad \alpha > \frac{1}{q}.$$

Then m is a Fourier multiplier of $L^p(\mathbb{R}^d)$ if $|1/p - 1/2| < 1/2q$.

Interpolating Theorem 1.5 with Corollary 1.4 (with p close to 1) yields

Corollary 1.6. *Let $m \in L^{\infty}(\mathbb{R}^d)$, $d \geq 2$, be homogeneous of degree zero and let $g_{\pm}$ be as in (1.5). Suppose that $1 < p < 4/3$ and*

$$\sup_{t \in (\mathbb{R}_+)^{d-1}} \|\mathcal{D}_1^{\alpha} \mathcal{D}_2^{\gamma} \cdots \mathcal{D}_{d-1}^{\gamma} [\beta_{(1)} \otimes \cdots \otimes \beta_{(d-1)} g_{\pm}(t_1, \dots, t_{d-1})]\|_{L^2(\mathbb{R}^{d-1})} < \infty, \quad \gamma > \frac{1}{2}, \quad \alpha > \frac{2}{p} - 1$$

and that the analogous conditions obtained by permuting the $(s_1, \dots, s_{d-1})$ -variables hold. Then m is a Fourier multiplier of $L^p(\mathbb{R}^d)$.

In particular if $\sup_{t \in (\mathbb{R}_+)^{d-1}} \|\beta_{(1)} \otimes \cdots \otimes \beta_{(d-1)} g_{\pm}(t_1, \dots, t_{d-1})\|_{\mathcal{H}_{\alpha}^2(\mathbb{R}^{d-1})} < \infty$ and $1 < p < 4/3$ then m is a Fourier multiplier of L^p in $\alpha > \frac{2}{p} - 1$. This result is essentially sharp: in §3 we show that in order for $\sup_{t \in (\mathbb{R}_+)^{d-1}} \|\beta_{(1)} \otimes \cdots \otimes \beta_{(d-1)} g_{\pm}(t_1, \dots, t_{d-1})\|_{\mathcal{H}_{\alpha}^q(\mathbb{R}^{d-1})} < \infty$ to imply that m is a Fourier multiplier of L^p we must necessarily have $\alpha \geq 2/p - 3/2 + 1/q$ if $1 < p < 4/3$ and $\alpha > 1/q$ if $4/3 \leq p \leq 2$.

In order to prove more refined results on $L^p(\mathbb{R}^d)$, $d \geq 3$, p close to 1, we shall use multiparameter Calderón-Zygmund theory. It turns out that it is useful (and easier) to first prove a result for the multiparameter Hardy-space $H^p(\mathbb{R}^d)$, $0 < p \leq 1$. The Hardy space H^p is defined in terms of square-functions invariant under the multiparameter family of dilations $\delta_t x = (t_1 x_1, \dots, t_d x_d)$, $t \in (\mathbb{R}_+)^d$. Again we formulate the multiplier result using localized multiparameter Sobolev spaces invariant under multiparameter dilations. In order to include a sharp result also for $p < 1$ we want to admit values of $q \leq 1$ in (1.2). To make this possible the definition of $\mathcal{H}_{\alpha}^q$ has to be modified. We may always assume that β above is such that $\sum_{r \in \mathbb{Z}} \beta^2(2^{-r} s) = 1$ for $s \neq 0$. Let $\psi_r = \beta^2(2^{-r} \cdot)$ if $r \geq 1$ and $\psi_0 = 1 - \sum_{r > 0} \psi_r$. For $n = (n_1, \dots, n_{d-1})$, $n_i \geq 0$, $i = 1, \dots, d-1$ set $\psi_n(\xi_1, \dots, \xi_{d-1}) = \prod_{i=1}^{d-1} \psi_{n_i}(\xi_i)$. The decomposition

$$g = \sum_{n \in (\mathbb{N}_0)^{d-1}} \hat{\psi}_n * g$$

is referred to as the inhomogeneous Littlewood-Paley decomposition of $\mathbb{R}^{d-1}$. Then

$$(1.9) \quad \|g\|_{\mathcal{H}_\alpha^q(\mathbb{R}^{d-1})} \approx \left\| \left(\sum_{n \in (\mathbb{N}_0)^{d-1}} 2^{2(n_1 + \dots + n_{d-1})\alpha} |\widehat{\psi}_n * g|^2 \right)^{1/2} \right\|_{L^q(\mathbb{R}^{d-1})}$$

for $1 < q < \infty$, and for $q \leq 1$ we define $\mathcal{H}_\alpha^q(\mathbb{R}^{d-1})$ as the space of tempered distributions for which the quasinorm on the right hand side of (1.9) is finite. In this paper we shall always have $\alpha > 1/q$; in this case $\mathcal{H}_\alpha^q$ is embedded in L^∞ . This and other properties of the spaces $\mathcal{H}_\alpha^q$ may be proved by obvious modifications of the one-parameter case; for the latter we refer to [27].

Theorem 1.7. *Let $m \in L^\infty(\mathbb{R}^d)$ be homogeneous of degree zero and related to $g_\pm$ as in (1.5). Suppose that $0 < r \leq 1$ and*

$$(1.10) \quad \sup_{t \in (\mathbb{R}_+)^{d-1}} \|\beta_{(1)} \otimes \dots \otimes \beta_{(d-1)} g_\pm(t_1 \cdot, \dots, t_{d-1} \cdot)\|_{\mathcal{H}_\alpha^r(\mathbb{R}^{d-1})} < \infty, \quad \alpha > \frac{2}{r} - 1.$$

Moreover if $d \geq 3$ suppose that

$$(1.11) \quad \sup_{t \in (\mathbb{R}_+)^{d-1}} \|\mathcal{D}_1^{2\gamma} \mathcal{D}_2^\gamma \dots \mathcal{D}_{d-2}^\gamma [\beta_{(1)} \otimes \dots \otimes \beta_{(d-1)} g_\pm(t_1 \cdot, \dots, t_{d-1} \cdot)]\|_{L^2(\mathbb{R}^{d-1})} < \infty, \quad \gamma > \frac{1}{r} - \frac{1}{2}$$

and that the analogous conditions obtained by permuting the $(s_1, \dots, s_{d-1})$ -variables hold. Then m is a Fourier multiplier of the multiparameter Hardy space $H^p(\mathbb{R}^d)$, $r \leq p < \infty$.

Note that in two dimensions Theorem 1.7 is a natural extension of Corollary 1.4 to H^p -spaces in product domains. The examples in §3 show that in higher dimensions additional assumptions such as (1.11) are necessary. When $d \geq 3$, Theorem 1.7 with $r = 1$ serves as a substitute for Theorem 1.1. Notice that if $r = 1$ condition (1.10) involves mixed derivatives in L^1 of order $d - 1 + \epsilon$, and condition (1.11) involves derivatives in L^2 up to order $(d - 1 + \epsilon)/2$. In comparison the hypotheses in Corollaries 1.3 and 1.6 involve L^2 derivatives up to order $(d + \epsilon)/2$ if p is close to 1. As a consequence we obtain the following analogue of Corollary 1.4, formulated in terms of the standard *oneparameter* Sobolev space L_α^q .

Corollary 1.8. *Let $m \in L^\infty(\mathbb{R}^d)$ be homogeneous of degree zero and related to $g_\pm$ by (1.5). Suppose that $q > 1$ and that*

$$\sup_{t \in (\mathbb{R}_+)^{d-1}} \|\beta_{(1)} \otimes \dots \otimes \beta_{(d-1)} g_\pm(t_1 \cdot, \dots, t_{d-1} \cdot)\|_{L_\alpha^q(\mathbb{R}^{d-1})} < \infty, \quad \alpha > \frac{d-1}{q}.$$

Then m is a Fourier multiplier of $L^p(\mathbb{R}^2)$ if $|1/p - 1/2| < 1/2q$.

The counterexamples in [22], [7] show that the statement of the Corollary is false in the range $|1/p - 1/2| > 1/2q$. However in view of Theorems 1.5 and 1.7 one expects the following sharper result. Namely suppose that for some $q \in (1, 2]$

$$(1.12) \quad \sup_{t \in (\mathbb{R}_+)^{d-1}} \|\beta_{(1)} \otimes \dots \otimes \beta_{(d-1)} g_\pm(t_1 \cdot, \dots, t_{d-1} \cdot)\|_{\mathcal{H}_\alpha^q(\mathbb{R}^{d-1})} < \infty, \quad \alpha > \frac{1}{q},$$

and in dimension $d \geq 3$ suppose that

$$(1.13) \quad \sup_{t \in (\mathbb{R}_+)^{d-1}} \|\mathcal{D}_1^\alpha \mathcal{D}_2^\gamma \dots \mathcal{D}_{d-2}^\gamma [\beta_{(1)} \otimes \dots \otimes \beta_{(d-1)} g_\pm(t_1 \cdot, \dots, t_{d-1} \cdot)]\|_{L^2(\mathbb{R}^{d-1})} < \infty, \quad \gamma > \frac{1}{2}, \quad \alpha > \frac{1}{q}$$

as well as the analogous conditions obtained by permuting the $(s_1, \dots, s_{d-1})$ -variables. Then m should be a Fourier multiplier of $L^p(\mathbb{R}^d)$ if $|1/p - 1/2| < 1/2q$. In order to prove this one is tempted to use analytic interpolation and interpolate between the L^{p_0} -estimate of Theorem 1.7, for p_0 close to 1, and the $L^{4/3}$ -estimate of Theorem 1.5. One would have to find the intermediate spaces for intersections of L^2 and L^q Sobolev spaces. However the intersection of the intermediate spaces does not need to be contained in the intermediate space of the intersections (for related counterexamples see [26]). It is actually possible to prove the result for $|1/p - 1/2| < 1/2q$ (assuming (1.12), (1.13)) by another approach. One has to use a general theorem for analytic families of operators acting on various kinds of atoms the proof of which relies heavily on multiparameter Calderón-Zygmund theory. We do not include the technical proof here but refer the reader to [5].

The paper is organized as follows: In §2 we prove Theorem 1.1 using weighted norm inequalities and variants of the maximal operator with respect to lacunary directions. Examples demonstrating the sharpness of our results in higher dimensions are discussed in §3. The proof of Theorem 1.5 is in §4; it relies on weighted norm inequalities which involve variants of the Kakeya maximal function. In §5 we prove the Hardy space estimates of Theorem 1.7.

As a convention we shall refer to the quasi-norms in H^p and $\mathcal{H}_\alpha^p$ as “norms” although for $p < 1$ these spaces are not normed spaces. By M_p , $1 \leq p \leq \infty$, we denote the standard space of Fourier multipliers of L^p . It will always be assumed that the even function $\beta \in C_0^\infty$ defined above satisfies $\sum_{\kappa \in \mathbb{Z}} [\beta(2^\kappa s)]^2 = 1$ for $s \neq 0$. If $\mathfrak{a} \in \{1, \dots, d\}$ and $k, \tilde{k}$ in $\mathbb{R}^{\mathfrak{a}}$ then we shall use the notation $k \leq \tilde{k}$ if $k_i \leq \tilde{k}_i$ for all $i \in \mathfrak{a}$. Similarly define $k \leq \tilde{k}$ etc. C will always be an abstract constant which may assume different values in different lines.

2. L^p -estimates in the plane

In the proof of Theorem 1.1 there is no loss of generality in assuming that m is supported in the quadrant where $\xi_1 > 0$ and $\xi_2 > 0$. By a limiting argument as in Stein’s book [25, p.109], it suffices to prove the theorem under the formally stronger assumption

$$\|h\|_\infty + \sup_{k \in \mathbb{Z}} \int_{I_k} |h'(s)| ds \leq A.$$

Let β be the smooth bump function defined in the introduction (supported in $\pm[5/8, 8/5]$). Let $\gamma \in C^\infty(\mathbb{R}^2 \setminus \{0\})$ be homogeneous of degree 1 such that $\gamma(\xi) = 1$ if $|\xi_1/\xi_2| \in [25/64, 64/25]$ (in particular on the support of $\beta \otimes \beta$) and such that $\gamma(\xi) = 0$ if $|\xi_1/\xi_2| \notin (1/4, 4)$. Set

$$h_\kappa(\xi) = \gamma(\xi) h(2^{-\kappa} \xi_1 / \xi_2).$$

Then we may split

$$m = \sum_{k \in \mathbb{Z}^2} [\beta \otimes \beta m_k](2^{k_1}, 2^{k_2}.)$$

where

$$\begin{aligned} m_k(\xi) &= \beta(\xi_1) \beta(\xi_2) h_{k_1 - k_2}(\xi_1 / \xi_2) \\ &= \beta(\xi_1) \beta(\xi_2) \int_0^{\xi_1 / \xi_2} h'_{k_1 - k_2}(s) ds. \end{aligned} \tag{2.1}$$

and h_κ is supported in $(1/4, 4)$, for all $\kappa \in \mathbb{Z}$. Also set

$$\widehat{T_k f}(\xi) = [\beta \otimes \beta m_k](2^{k_1} \xi_1, 2^{k_2} \xi_2) \widehat{f}(\xi).$$

Then by standard multiparameter Littlewood-Paley theory and duality, to establish Theorem 1.1 for $p \in [2, p_0)$, $p_0 < \infty$, it suffices to obtain an inequality

$$(2.2) \quad \int |T_k f|^2 \omega \leq C A^2 \int |f|^2 \mathfrak{M} \omega$$

for a certain operator $\omega \mapsto \mathfrak{M} \omega$ which is bounded on $L^q(\mathbb{R}^2)$ for $(p_0/2)' < q \leq \infty$. By our assumption on h ,

$$(2.3) \quad \sup_{\kappa \in \mathbb{Z}} \int |h'_\kappa(s)| ds \leq C A.$$

We denote by $\mathcal{L}_k$ the standard Littlewood-Paley operator, such that

$$\widehat{\mathcal{L}_k f}(\xi) = \beta(2^{k_1} \xi_1) \beta(2^{k_2} \xi_2) \widehat{f}(\xi)$$

and define the operator $S_{k,s}$ by

$$\widehat{S_{k,s} f}(\xi) = \begin{cases} \widehat{f}(\xi), & \text{if } 2^\kappa \xi_1 / \xi_2 > s, \xi_1 \geq 0, \xi_2 \geq 0 \\ 0, & \text{otherwise} \end{cases}.$$

Then from (2.1) we see that

$$T_k f(x) = \int_{1/8}^8 \mathcal{L}_k S_{k_1-k_2,s} f(x) h'_{k_1-k_2}(s) ds$$

Then, if $\omega \geq 0$ is a weight we apply the Cauchy-Schwarz inequality to obtain

$$(2.4) \quad \int |T_k f(x)|^2 \omega(x) dx \leq C A \iint |\mathcal{L}_k S_{k_1-k_2,s} f(x)|^2 |h'_{k_1-k_2}(s)| ds \omega(x) dx.$$

Let $M_{(1)}$, $M_{(2)}$ be the Hardy-Littlewood maximal functions with respect to the coordinate directions and let $M_{\kappa,s}$ be the Hardy-Littlewood maximal function with respect to the direction perpendicular to $\{\xi; 2^\kappa \xi_1 / \xi_2 = s\}$, i.e. in the direction $(1, -2^{-\kappa} s)$. Then using weighted norm inequalities for singular integral operators due to Córdoba and Fefferman ([13], see also [18]) we see that the expression on the right hand side of (2.4) is dominated by

$$C_\alpha A \int |f(x)|^2 M_{(1)} M_{(2)} \left[\int_{1/8}^8 (M_{k_1-k_2,s} \omega^\alpha)^{1/\alpha} |h'_{k_1-k_2}(s)| ds \right] (x) dx$$

where $\alpha > 1$. Now the proof of (2.2) is completed by the following

Proposition 2.1. *Let, for $\alpha \geq 1$,*

$$\mathfrak{M}_\alpha \omega(x) = \sup_{\kappa \in \mathbb{Z}} \int_I (M_{\kappa,s} \omega^\alpha)^{1/\alpha}(x) |\lambda_\kappa(s)| ds$$

where $I = [1/8, 8]$ and

$$\sup_{\kappa \in \mathbb{Z}} \int_I |\lambda_\kappa(s)| ds \leq B < \infty.$$

Then $\mathfrak{M}_\alpha$ is bounded on $L^p(\mathbb{R}^2)$, $\alpha < p < \infty$, with norm $\leq C_{p,\alpha} B$.

Proof. Since

$$\mathfrak{M}_\alpha(\omega) \leq B^{1-\frac{1}{\alpha}} [\mathfrak{M}_1(\omega^\alpha)]^{\frac{1}{\alpha}}$$

it suffices to prove that $\mathfrak{M}_1$ is bounded on L^p , $1 < p < \infty$ with norm $C_p B$. If $\alpha > 1$ then $\mathfrak{M}_\alpha$ will be bounded on L^p , $p > \alpha$, with norm $\leq C_{p/\alpha} B$.

We follow arguments by Nagel, Stein and Wainger [23] as modified by Christ (see [2]). Let $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ be smooth, even, nonnegative, with $\varphi(0) > 0$ such that $\widehat{\varphi}$ has compact support in $[-1/20, 1/20]$. Let

$$\psi(\xi_1, \xi_2) = \widehat{\varphi}(\xi_1 + \xi_2)$$

and define for $\kappa \in \mathbb{Z}$

$$\widehat{P_{\kappa s}^l \omega}(\xi) = \psi(2^l \xi_1, s 2^{l-\kappa} \xi_2) \widehat{\omega}(\xi).$$

It suffices to show that for $1 < p < \infty$, N being an arbitrary positive integer

$$(2.5) \quad \left\| \sup_{-N \leq \kappa \leq N} \int_I \sup_{l \in \mathbb{Z}} |P_{\kappa s}^l \omega| |\lambda_\kappa(s)| ds \right\|_p \leq C_p B \|\omega\|_p$$

where C_p is independent of N . Then an application of the monotone convergence theorem allows to pass to the limit. We note that for fixed κ , and $1 < p \leq \infty$

$$(2.6) \quad \begin{aligned} \left\| \int_I \sup_{l \in \mathbb{Z}} |P_{\kappa s}^l \omega| |\lambda_\kappa(s)| ds \right\|_p &\leq C \int_I \left\| \sup_{l \in \mathbb{Z}} |P_{\kappa s}^l \omega| \right\|_p |\lambda_\kappa(s)| ds \\ &\leq C_p B \|\omega\|_p \end{aligned}$$

by the L^p estimate for the one-dimensional Hardy-Littlewood maximal function $M_{k,s}$. This means that we know a priori that the left hand side of (2.5) is bounded by $B C_p(N) \|f\|_p$ (with $C_p(N) \leq C'_p N$) and it remains to be shown that $C_p(N)$ can be chosen independently of N . In what follows we define $C_p(N)$ to be the best constant in (2.5).

We first consider the case $2 \leq p < \infty$. Since the L^∞ -estimate is trivial it suffices to prove the L^2 inequality. We smoothly split ψ into two parts, $\psi = \psi^0 + \psi^1$ with ψ^1 supported in the unit ball and ψ^0 supported in the cone $\{\xi; |\xi_1 + \xi_2|/|\xi| \leq 1/2\}$. We correspondingly define the operators $P_{\kappa s}^{l,0}$ and $P_{\kappa s}^{l,1}$. Note that there is the pointwise inequality

$$(2.7) \quad |P_{\kappa s}^{l,1} \omega(x)| \leq C M_{(1)} M_{(2)} \omega(x)$$

which implies

$$(2.8) \quad \left\| \sup_{\kappa} \int_I \sup_{l \in \mathbb{Z}} |P_{\kappa s}^{l,1} \omega| |\lambda_\kappa(s)| ds \right\|_p \leq C_p B \|\omega\|_p, \quad 1 < p \leq \infty.$$

Concerning $P_{\kappa s}^{l,0}$ we have

$$|P_{\kappa s}^{l,0} \omega(x)| \leq C [M_{(1)} M_{(2)} \omega(x) + M_{\kappa,s} \omega(x)]$$